

FEAST[®]

...Clearly the Liquid Leader

Corn and Small Grains

ROW PLACEMENT GUIDELINES

Based On Cation Exchange Capacity and Percent Organic Matter

Row placement (in-row with direct seed contact) of starter fertilizers such as FEAST[®] Yield Master[™] 9-18-9, 3-18-18, 8-16-11w2S, 3-15-19w3S, and FEAST[®] Side-Kick[™] 0-0-25 w17S should be avoided in soil conditions that are low in cation exchange capacity and/or low in organic matter. Conklin recommends an AgroVantage[®] soil test through Midwest Labs, Omaha, Nebraska, to accurately determine these levels. The data has been calibrated only to Conklin fertilizer products and should be interpreted and used as such. Refer to the product specification sheets for additional information.

STATEMENT 1

No row placement of fertilizer should be made on soils having C.E.C. of less than 4.0. For soils having C.E.C less than 4.0, fertilizer must be placed without direct seed contact such as with a splitter.

STATEMENT 2

No row placement of fertilizer should be made on soils having an organic matter content of less than 0.5 percent.

STATEMENT 3

For the C.E.C. values 4-13, row application should only be made if the organic matter is equal to or exceeds those values listed.

If Cation Exchange Capacity is:

The Organic Matter Percent should be:

Equal to or greater than 4.....	Equal to or greater than 4.50
Equal to or greater than	Equal to or greater than 4.05
Equal to or greater than 6.....	Equal to or greater than 3.60
Equal to or greater than 7.....	Equal to or greater than 3.15
Equal to or greater than 8.....	Equal to or greater than 2.70
Equal to or greater than 9.....	Equal to or greater than 2.25
Equal to or greater than 10	Equal to or greater than 1.80
Equal to or greater than 11.....	Equal to or greater than 1.5
Equal to or greater than 12.....	Equal to or greater than 0.90
Equal to or greater than 13.....	Equal to or greater than 0.5

STATEMENT 4

Soils outside of these guidelines should be made using in-row, non-seed contact application such as a Y-Not splitter or 2x2.

STATEMENT 5

Total application not to exceed 5 gallons of Feast NPK Starter plus 1 gallon of Side-Kick per acre according to these guidelines. 8-16-11w2S and 3-15-19w3S may be applied up to 6 gallons per acre, but should not be applied with additional Side-Kick 0-0-25w17S.

